

AD-A051 759

FLORIDA UNIV GAINESVILLE DEPT OF INDUSTRIAL AND SYS--ETC F/G 12/1  
OBSERVATIONS ON THE MINIMUM SPHERE PROBLEM.(U)

MAY 77 D HEARN

DAHC04-75-6-0150

UNCLASSIFIED

RR-77-5


ARO-12640.26-M

NL

| OF |  
AD  
A061759



END  
DATE  
FILMED  
4-78  
DDC



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

AKD 12640.26-11

AD A051759

4

# RESEARCH REPORT

AD NO. \_\_\_\_\_  
DDC FILE COPY

DDC  
RECEIVED  
MAR 24 1978  
B

Industrial & Systems  
Engineering Department  
University of Florida  
Gainesville, FL 32611

**DISTRIBUTION STATEMENT A**

Approved for public release;  
Distribution Unlimited

4

OBSERVATIONS ON THE MINIMUM  
SPHERE PROBLEM

Research Report No. 77-5

by

Donald Hearn

May 1977

Department of Industrial and Systems Engineering  
University of Florida  
Gainesville, Florida 32611

APPROVED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED

This research was supported in part by the Army Research Office,  
Triangle Park, NC, under contract number DAHC04-75-G-0150, and  
NSF contract number ENG74-22159.

DDC  
RECEIVED  
MAR 24 1978  
B

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER (14) 12640, 26-M	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) (6) Observations on the Minimum Sphere Problem	5. TYPE OF REPORT & PERIOD COVERED (9) Research report Technical	
7. AUTHOR(s) (18) ARO	6. PERFORMING ORG. REPORT NUMBER (14) RR-77-57	
7. AUTHOR(s) (14) Donald Hearn	8. CONTRACT OR GRANT NUMBER(s) (15) DAHC04-75-G-0150 (16) -ENG74-22159	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Industrial and Systems Engineering University of Florida Gainesville, Florida 32611	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS (16) 20061102A14D Rsch in & Appl of Applied Math.	
11. CONTROLLING OFFICE NAME AND ADDRESS U.S. Army Research Office NSF P.O. Box 12211 Washington, DC Triangle Park, NC 27709 22550	12. REPORT DATE (11) May 1977	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	13. NUMBER OF PAGES (12) 22p 21	
	15. SECURITY CLASS. (of this report) Unclassified	
	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE N/A	
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)  N/A		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)  For a subset A of Euclidean n-space, the location problem of finding the point $x$ which minimizes the maximum distance $d(x, a)$ for $a$ in A may be interpreted as finding the smallest sphere which encloses A. Algorithms have previously been developed for the situations when A is finite or a polytope. Here we concentrate on interpretations and properties of the problem, particularly its relation to other problems: cases where the sphere problem is dual to that of finding a shortest vector in A, where there is a connection with a		

DD FORM 1 JAN 73 1473 EDITION OF 1 NOV 65 IS OBSOLETE

Unclassified  
SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

404 399

EB



## 20. Abstract (cont)

→ maximum moment-of-inertia problem, etc. Conjectures, relating the minimum sphere to points in A which define the diameter of A, are also discussed.

ACCESSION for	
NTIS	White Section <input checked="" type="checkbox"/>
DDC	Buff Section <input type="checkbox"/>
UNANNOUNCED	<input type="checkbox"/>
JUSTIFICATION _____	
BY _____	
DISTRIBUTION/AVAILABILITY CODES	
Dist. A, AIL, and/or SPECIAL	
A	

## TABLE OF CONTENTS

ABSTRACT . . . . .	1
SECTIONS	
Introduction . . . . .	1
Primal and Dual Formulation . . . . .	2
Duality Interpretations . . . . .	4
Solution Methods . . . . .	8
The Minimum Circle and Far Points . . . . .	13
REFERENCES . . . . .	16

# ABSTRACT

For a subset  $A$  of Euclidean  $n$ -space, the location problem of finding the point  $\underline{x}$  which minimizes the maximum distance  $d(\underline{x}, \underline{a})$  for  $\underline{a}$  in  $A$  may be interpreted as finding the smallest sphere which encloses  $A$ . Algorithms have previously been developed for the situations when  $A$  is finite or a polytope. Here we concentrate on interpretations and properties of the problem, particularly its relation to other problems: cases where the sphere problem is dual to that of finding a shortest vector in  $A$ , where there is a connection with a maximum moment-of-inertia problem, etc. Conjectures, relating the minimum sphere to points in  $A$  which define the diameter of  $A$ , are also discussed.



## 1. Introduction

For a subset  $S$  of  $E^n$ , the location problem of finding the point  $x$  which minimizes the maximum of  $\|x - a\|$  for  $a \in S$ , may be interpreted as finding the smallest sphere which encloses  $S$ . Previously, algorithms have been developed for solving this problem when  $S$  is finite or a polytope [4, 6]. In this note we concentrate on interpretations and properties of the problem; particularly its relation to other problems. For example, we show a case where the sphere problem is dual to the problem of finding the vector of minimum norm in  $S$ . As another example (in  $E^2$ ) we show the relation to a maximum moment of inertia problem. Other aspects discussed concern conjectures relating the minimum sphere to points in  $S$  which define the diameter of  $S$ .

## 2. Primal and Dual Formulation

The minimum sphere for a given finite set of points  $a_i \in E^n$ ,  $i = 1, 2, \dots, m$ , may be determined by solution of the mathematical program

$$(P1) \quad \min_{(r,c)} \quad r$$

$$\text{s.t.} \quad r \geq \|c - a_i\| \quad i = 1, 2, \dots, m$$

with variables  $r$  and  $c$ . The solution of (P1),  $(r^*, c^*)$ , gives the radius and center of the sphere (the existence and uniqueness of  $(r^*, c^*)$ , for any bounded set is guaranteed [1]). Problem (P1) is equivalent to

$$(P2) \quad \min \quad s$$

$$\text{s.t.} \quad s \geq \|c - a_i\|^2 \quad i = 1, 2, \dots, m$$

where  $s = r^2$ . By introducing the variable  $z = s - c^T c$  [9], (P2) becomes the convex quadratic program

$$(P3) \quad \min_{(z,c)} \quad z + c^T c$$

$$\text{s.t.} \quad z + 2a_i^T c \geq a_i^T a_i \quad i = 1, 2, \dots, m$$

The dual of (P3) is the concave quadratic program [4]:

$$(D3) \quad \max_v \quad h(v) = \sum v_i (a_i^T a_i) - v^T A^T A v$$

$$\text{s.t.} \quad \sum v_i = 1 \quad v_i \geq 0 \quad i = 1, 2, \dots, m$$

where  $A$  has columns  $a_i$ .

A short note by Kuhn [8] shows that these problems (P3) and (D3) are "hybrid" programs, i.e., they are linear programs with a sum of squares added to the objective function. (P3) has this form and so does (D3) if we define  $w \equiv Av$  so the second term in the objective is  $w^T w$ .

If  $v^*$  solves (D3) and  $(s^*, c^*)$  solves (P2) then

$$s^* = h(v^*)$$

$$\text{and} \quad c^* = \sum v_i^* a_i.$$

Note that the latter equation and the conditions on  $v$  in (D3) express the intuitively obvious fact that  $c^*$  is a convex combination of the  $a_i$ . The coefficients expressing the optimal combination are the Kuhn-Tucker multipliers which solve (D3). By a theorem of Caratheodory [10], the convex combination may always be expressed with at most  $n + 1$  positive coefficients. These facts are easily visualized for  $a_i \in E^2$ . The minimum circle will either be defined by the two most distance points, or it will pass through the vertices of three points which define a non obtuse triangle. Of course, more points may lie on the circle, but two or three suffice to express the center as a convex combination.

### 3. Duality Interpretations

In this section we show that two problems, one in mechanical physics and the other a geometrical "least distance" problem, are dual to the minimum sphere problem.

The first of these might be called a maximum moment of inertia problem. (I am grateful to my colleague, E. J. Muth, for pointing out this problem and its relation to the sphere problem.)

It is a basic principle of physics that a body set into rotation will tend to rotate about its center of gravity. This happens, for example, when a frisbee is sent spinning through the air. The physical principle involved is that natural forces tend to minimize the rotational energy of the body by choosing the center of gravity as the axis of rotation.

Consider now the following problem: Suppose we wish to assign masses  $m_i$  to a finite set of points  $a_i$  on a body so as to maximize its moment of inertia. We assume that the  $a_i$  are connected by thin rods or in some other manner involving negligible mass. For example, the  $a_i$  may be points on a thin lamina. Without loss of generality, we may assume the total mass to be distributed among the points is equal to one. Then, since the formula for moment of inertia,  $I$ , is

$$I = \sum m_i r_i^2$$

where  $r_i$  is the distance from the point  $a_i$  to the axis of rotation, we may write the problem as

$$\begin{aligned} \text{(PI)} \quad & \max_{m_i} \sum m_i ||a_i - \bar{x}||^2 \\ & \text{s.t.} \quad \sum m_i = 1 \quad m_i \geq 0 \quad i = 1, 2, \dots, m \end{aligned}$$

where  $\bar{x}$  is the center of gravity. But owing to the minimizing feature of the center of gravity we may write



$$\begin{aligned}
(\text{PI1}) \quad & \max_{m_i} \min_x \sum m_i ||a_i - x||^2 \\
& \text{s.t.} \quad \sum m_i = 1, \quad m_i \geq 0 \quad i = 1, 2, \dots, m
\end{aligned}$$

Thus the moment of inertia problem, (PI), is a maximin problem, and by a well known theorem [10, Corollary 37.3.2], the max min in (PI1) may be interchanged to min max. This yields

$$\begin{aligned}
(\text{DI1}) \quad & \min_x \max_{m_i} \sum m_i ||a_i - x||^2 \\
& \text{s.t.} \quad \sum m_i = 1 \quad m_i \geq 0 \quad i = 1, 2, \dots, m.
\end{aligned}$$

The inner maximization is easily solved by setting  $m_j = 1$ , where  $||a_j - x||^2 = \max_i ||a_i - x||^2$ , thus we obtain

$$(\text{DI2}) \quad \min_x \max_i ||a_i - x||^2$$

Which is equivalent to (P2) where  $c = x$  and  $c^* = \bar{x}$ . Thus we have proven that the solution of the problem (PI) is obtained by constructing the minimum circle for the given  $a_i$  and assigning the weights  $m_i = v_i^*$ , where  $v^*$  is the vector of optimal Kuhn-Tucker multipliers.

It is interesting that the theorem of Caratheodory quoted earlier states that, for this problem, the maximum number of points with positive weight need be no greater than three. Also, the angular kinetic energy and the radius of gyration of a rotating body are proportional to the moment of inertia, so maximizing these quantities yields the same result.

The interpretation given above does not extend directly for  $a_i \in E^3$  because the axis of rotation is a line rather than a point.

Our second problem arises in the minimization of nondifferentiable convex functions [13]. An important subproblem is to determine the vector,  $d^*$ , of minimum norm in a given compact convex set  $S$ , i.e., solve the problem:



$$\begin{aligned}
 \text{(PM)} \quad & \min ||d||^2 \\
 & \text{s.t. } d \in S.
 \end{aligned}$$

If we make the assumption that  $S$  is a polytope with extreme points  $a_i$ ,  $i = 1, 2, \dots, m$ , then we have

$$\begin{aligned}
 \text{(PM1)} \quad & ||d^*||^2 = \min_v ||\sum v_i a_i||^2 \\
 & \text{s.t. } \sum v_i = 1 \quad v_i \geq 0 \quad i = 1, 2, \dots, m.
 \end{aligned}$$

If we further assume that the  $a_i$  have the same Euclidean norm, say  $||a_i|| = 1$ , then from (PM1) we obtain

$$\begin{aligned}
 1 - ||d^*||^2 &= 1 - \min ||\sum v_i a_i||^2 \\
 &= 1 + \max - ||\sum v_i a_i||^2 \\
 &= \max 1 - ||\sum v_i a_i||^2 \\
 &= \max \sum v_i (a_i^T a_i) - ||\sum v_i a_i||^2 \\
 & \text{s.t. } \sum v_i = 1, v_i \geq 0.
 \end{aligned}$$

Comparison with (D3) gives the (Pythagorean) dual relation

$$||d^*||^2 + s^* = 1$$

where  $s^*$  is the radius squared of the minimum sphere covering the  $a_i$  and  $d^* = c^*$ .


See Figure 1 for a sketch. See [7] for computational experience where such normalizing of the  $a_i$  has proven helpful in nondifferentiable optimization.

It would be interesting to know whether there is a geometrical interpretation for the case when not all  $||a_i|| = 1$ .

As an aside on problem (PM1), we note that it may be written as

$$(PM2) \quad \max_d [ - d^T d + 2 \min_i a_i^T d ].$$

This formulation calls for the *unconstrained* maximization of a strictly concave function, but as far as we know, it has not been exploited in any solution procedure. To obtain (PM2) from (PM1), one has only to regard the latter as a quadratic programming problem and formulate Dorn's dual [3]. We omit the details of the derivation.



Note:  $(r^*)^2 = s^*$

$$||d^*||^2 + (r^*)^2 = 1$$

FIGURE 1

#### 4. Solution Methods

Since both (P3) and (D3) are quadratic programs, either could be solved by one of the well known algorithms designed for such problems. Care should be taken, however, because both objectives are *semidefinite*. Thus methods such as the Theil-Van de Panne procedure [12] which require a *definite* quadratic objective are not directly applicable, but others, such as Dantzig's procedure [2] are. If  $m$  is small, then (P3) or (D3) could both be solved with about the same effort. If, however,  $m$  is very large, (D3) is more attractive because there is only one constraint. In [4] Elzinga and Hearn give a decomposable solution procedure for (D3) which uses the Dantzig algorithm and for which the size of the tableau is independent of  $m$ . The basis for the decomposition is Caratheodory's theorem. A similar algorithm for the case when  $S$  is a polytope is described in [6].

For the problem in the plane one could also employ quadratic programming, but the following geometrical procedure quoted from [5] is more efficient:


- "1. Pick any two given points and go to step 2.
2. Let the two points define the diameter of a circle. If this circle covers all points, stop. Otherwise choose some point outside the circle and the two defining points and go to step 3.
3. If the three points define a right triangle or an obtuse triangle, drop the point at the angle  $\geq 90^\circ$  and go to step 2 with the remaining two points. Otherwise if the triangle has strictly acute angles, go to step 4.
4. If the circle defined by the three points covers all points, stop. Otherwise choose some outside point, call it  $\sigma$ , and label as  $\alpha$  a point from among the three defining points that is farthest from  $\sigma$ . Extend the diameter of the current circle through point  $\alpha$  to divide

the plane into two half-planes. Label as  $\beta$  the defining point in the same half-plane with  $\sigma$  and as  $\gamma$  the remaining point. With points  $\alpha$ ,  $\gamma$ , and  $\sigma$ , go to step 3.

In step 3 if the three points are on a straight line, we consider this a 'triangle' with a  $180^\circ$  angle at the intermediate point. Also note that by considering a right triangle as a two-point case we assume that in step 4 the point  $\beta$  is strictly contained in the half-plane with  $\sigma$ . In practice it seems reasonable to choose the farthest point outside the current circle rather than just some point as indicated in steps 2 and 3.

Convergence of this algorithm is based on showing that the circles created in each iteration are monotone increasing in radius. Then since there are only finitely many two-point and three-point circles, the process is finite."

We quote the algorithm here (but not the convergence proof) to emphasize (a) that it is initiated with an arbitrary pair of points, and (b) that once a point is covered, it need not remain covered (see Figure 2). All the procedure guarantees is that successive circles strictly increase in radius.



Point  $a_1$  uncovered by third circle.

FIGURE 2



The efficiency of this geometrical algorithm, as measured by empirical testing, appears to be good for  $m \leq 100$ . (See Figures 3 and 4.) For large  $m$ , however, the time is probably of order  $m^2$ ; in fact, this is suggested by Figure 4.

Whether a more efficient method can be devised is an open question. Certainly, an algorithm which kept all prior points covered would be of order  $m^*$ , but would require that each circle be *efficiently* constructed from the previous one. In other words, the critical question would be how to construct the minimum circle for  $k + 1$  points, knowing the minimum circle for  $k$  points. The answer is trivial for  $k + 1 = 2$  or  $3$ , but for  $k + 1 = 4$ , there are  $\binom{4}{3} + \binom{4}{2} = 10$  possible circles to consider, and for larger  $k$  the subproblem seems no easier than the original.

---

\* Shamos and Hoey [11] have conjectured that an algorithm of order  $m \log_2 m$  is possible.






Figure 3





Figure 4

### 5. The Minimum Circle and Far Points

For a finite point set  $S = \{a_1, a_2, \dots, a_m\}$ ,  $a_i \in E^2$ , we define as *far points* any pair  $(a_k, a_\ell)$  for which  $\|a_k - a_\ell\| = \max_{i,j} \|a_i - a_j\| = \text{diameter of } S$ .

It is natural to conjecture that the minimum circle passes through at least one far point. Figure 5 represents a counterexample. The far points are  $a_1$  and  $a_2$  because the two arcs, Arc 1 and Arc 2 enclose all  $a_i$ . Clearly, however, the minimum circle is defined by  $a_3, a_4$ , and  $a_5$ . (The distances from  $a_5$  to  $a_3$  and  $a_4$  are less than the diameter of the circle.)



Minimum Circle defined by  
 $\{a_3, a_4, a_5\}$ .

FIGURE 5

The same figure enables us to determine the probability that a circle defined by two points will enclose a random third point.

Theorem Let  $a_1$  and  $a_2$  be fixed points and assume a random third point is no farther from  $a_1$  and  $a_2$  than  $||a_1 - a_2||$ . Then the probability  $P$ , that the circle defined by  $a_1$  and  $a_2$  covers the third point is

$$P = \frac{3\pi}{8 - 6\sqrt{3}} \doteq .64.$$

Proof: Consider Figure 6 where  $a_1$  and  $a_2$  (distance  $d$  apart) represent the two fixed points so that the third is enclosed in the area bordered by Arc 1 and Arc 2. The probability of it not being exterior to the circle defined by  $a_1$  and  $a_2$  is simply the ratio of the area of the circle to the area between the arcs.

Point  $a_2$  is placed at the origin and the arcs each have a radius of  $d$ . We determine the area between the arcs by integration over one-quarter of the area (shaded portion):

$$\text{Area between arcs} = 4 \int_0^{\bar{x}} ((d^2 - x^2)^{1/2} - \frac{d}{2}) dx,$$

where  $\bar{x}$  is determined by

$$\frac{d^2}{4} + \bar{x}^2 = d^2 \Rightarrow \bar{x} = \pm \frac{d}{2} \sqrt{3}.$$

So,

$$\begin{aligned} \text{area between arcs} &= 4 \int_0^{\frac{d}{2} \sqrt{3}} ((d^2 - x^2)^{1/2} - \frac{d}{2}) dx \\ &= 2 \left[ x \sqrt{d^2 - x^2} + d^2 \sin^{-1} \frac{x}{d} \right] \frac{d}{2} \sqrt{3} \\ &\quad - 4d \left[ \frac{x}{2} \right]_0^{\frac{d}{2} \sqrt{3}} \\ &= 2 \left( \left( \frac{d}{2} \sqrt{3} \right) \sqrt{d^2 - \frac{3d^2}{4}} + d^2 \sin^{-1} \frac{\sqrt{3}}{2} \right) - d^2 \sqrt{3} \end{aligned}$$



$$= 2d^2 \left( \sqrt{\frac{3}{4}} + \sin^{-1} \sqrt{\frac{3}{2}} - \sqrt{\frac{3}{2}} \right)$$

$$= 2d^2 \left( \frac{\pi}{3} - \sqrt{\frac{3}{4}} \right).$$

Since the area of the circle is  $\pi \frac{d^2}{4}$ , the ratio is

$$P = \frac{\pi}{8 \left( \frac{\pi}{3} - \sqrt{\frac{3}{4}} \right)} = \frac{3/2 \pi}{4\pi - 3\sqrt{3}} = \frac{3\pi}{8\pi - 6\sqrt{3}} \doteq .64.$$




FIGURE 6

This result suggests further problems of the same sort. For example, what is the probability that three points uniformly distributed over some convex set in  $R^2$  will form a triangle with an obtuse angle (so that the minimum circle is defined by the two far points)? When the set is the unit square, Monte Carlo simulation shows that the answer is approximately .70. To our knowledge, the question posed has not been answered analytically. For the interested researcher, the monograph by Kendall and Moran [14] is a starting point.



## References

1. Blumenthal, L. M. and Wahlin, G. E., "On the Spherical Surface of Smallest Radius Enclosing a Bounded Subset of  $n$  - Dimensional Euclidean Space," American Mathematical Society Bulletin, Vol. 47 (1941).
2. Dantzig, G. B., Linear Programming and Extensions, Princeton University Press (1963).
3. Dorn, W. S., "Duality in Quadratic Programming," Quarterly of Applied Mathematics, 18 (1960).
4. Elzinga, J. and Hearn, D. W., "The Minimum Covering Sphere Problem," Management Science 19, 96-104 (1972).
5. Elzinga, J. and Hearn, D. W., "Geometrical Solutions for Some Minimax Location Problems," Transportation Science, Vol. 6, No. 4 (1972).
6. Elzinga, J. and Hearn, D. W., "The Minimum Sphere Covering a Convex Polyhedron," NRLQ, Vol. 21, No. 4 (1974).
7. Hearn, D. W. and Lowe, T. J., "A Subgradient Procedure for the Solution of Minimax Location Problems," to appear in Computers and Industrial Engineering.
8. Kendall, M. G. and Moran, P. A. P., "Geometrical Probability," Griffin's Statistical Monograph No. 10, Hafner Publishing Co., New York, 1963.
9. Kuhn, H. W., "Sylvester's Problem is a Hybrid Program," unpublished note, Princeton University, March 1975.
10. Nair, K. P. K. and Chandrasekaran, R., "Optimal Location of a Single Service Center of Certain Types," NRLQ, 18 (1971).
11. Rockafellar, R. T., Convex Analysis, Princeton University Press, 1970.
12. Shamos, M. I. and Hoey, D. "Closest Point Problems," Proceedings 16th Annual Symposium on Foundations of Computer Science. (IEEE), Berkeley, Calif. (1975).
13. Theil, H. and van de Panne, C., "Quadratic Programming as an Extension of Conventional Quadratic Maximization," Management Science, Vol. 7 (1960).
14. Wolfe, P., "Algorithm for a Least-Distance Programming Problem," Mathematical Programming Study 1 (1974).